


In the case of fever (temperature above 38 °C), swelling, pain and redness at the wound site or abdominal pain with bowel movement complaints (constipation or diarrhoea), you should visit a doctor or our outpatient clinic without delay.

We hope to have answered all your potential questions with this information and wish you a speedy recovery.

Your surgical team

Surgery

Discharge information



Behaviour after endoscopic removal of the gall bladder

Publication details:

Media owner: NÖ Landesgesundheitsagentur

Version 05/2015

Galle

Dear Patient,

Further to our discussion at discharge and the doctor's letter, we would like to provide you with some more information relating to the time after your stay in our department.

Please observe the following for the most favourable treatment outcomes:

Physical rest	2-4 weeks
Avoid lifting and carrying heavy objects (3–5 kg)	2-4 weeks
Sporting activities, full physical activity after	2-4 weeks
Removal of stitches, staples or Steri-Strips	Depending on the medical orders 7-10 days after surgery
In case of souble stitches, removal of the plaster by the patient	Approx. one week after surgery

Important general information

Nutrition

We recommend a light normal diet for the first two to four weeks after surgery. After that you can, in principle, eat and drink anything you like.

However, initially you should only eat smaller portions of high fat foods, so that your body can gradually become accustomed to the new situation.

Postoperative pain

Pain may occur in the area around the surgical wound, however this usually subsides within one week.

A slightly swollen abdomen with shooting pains radiating into the shoulder is possible in the first week after surgery; this is due to the gas used with the surgical technique.

Personal hygiene

Showering is possible from the second day after surgery. However, the wound area should only be cleaned with water, and no soap. Please discuss the use of waterproof bandages with the doctor treating you. Full baths should be taken at the earliest after the stitches are removed.

In principle, no specific scar treatment is required; however regularly applying a skin care cream can greatly aid the healing process.