


In the case of atypical changes in the affected area, fever, bleeding and wound infection, you should visit our specialist department/outpatient clinic without delay.

We hope to have answered all your potential questions with this information and wish your son a speedy recovery.

Your treatment team

Publication details:
Media owner: NÖ Landesgesundheitsagentur
Version 05/2015
Vorhaut OP

Discharge information



Behaviour after foreskin surgery

Behaviour after foreskin surgery ■

Dear parents,

Dear patient,

A foreskin operation has been carried out on you/your son.

Further to our discussion at discharge and the doctor's letter, we would like to provide you with some more information relating to the time after your stay in our department.

For the best possible healing, we ask that you adhere to the following patterns of behaviour:

Behaviour after foreskin surgery ■

- Physical rest (no physical education classes) for one to two weeks
- In the first three days, the wound area should only be cleaned with a damp flannel (camomile tea)
- Full baths from the 10th day (otherwise the suture material dissolves too soon)
- Swelling of the surgical wound is normal in the early days and can be treated with painkillers and/or anti-inflammatory ointments.

Behaviour after foreskin surgery ■

- To prevent readhesion during foreskin preservation surgery, the foreskin should be pulled forward and then pushed back regularly.
- Any whitish/yellowish coatings may be caused by already dissolved stitches.

A check-up by the consultant is intended, depending on the medical prescription.